عنوان طرح تحقیقاتی: بررسی عوامل مرتبط بر ارتقای رفتارهای پیشگیری کننده از پدیکولوزیس دردانش آموزان دختر مقطع ابتدایی شهرتربت حیدریه: کاربردی از مدل باور سلامتی

تاریخ خاتمه طرح :1403/08/20

مجری یا محقق اصلی و همکاران با ذکر وابستگی هر فرد:
دکتر هادی علیزاده سیوکی عضو هیات علمی گروه بهداشت عمومی

عنوان پیام پژوهشی (حداکثر 20 کلمه): به نظرمی رسد شیوع آلودگی به شپش سر در دانش آموزان مدارس ابتدایی خصوصا

کشورهای درحال توسعه رو به افزایش است

پیام کلیدی (حداکثر80 کلمه) : : حساسیت درک شده و شدت درک شده نقش بسزایی در ارتقاء رفتارهای پیشگیری کننده دارد.

متن پیام پژوهشی (حداکثر240 کلمه): وضعیت نامطلوب اقتصادی و اجتماعی، تراکم جمعیت، پایین بودن سطح استاندارد زندگی، فقر بهداشتی و جنسیت در اشاعه ی آلودگی به شپش سر مؤثر می باشد . مدارس در عین داشتن پتانسیل بالا جهت شیوع آلودگی، مناسب ترین مکان جهت ارایه آموزش های بهداشتی لازم و در نتیجه پیشگیری و کنترل آلودگی به شپش سر هستند

- اهمیت موضوع(60کلمه)،
- مهمترین نتایج طرح به زبان غیر تخصصی(100 کلمه)
- موارد کاربرد نتایج طرح (80 کلمه)

تأثیرات و کاربردها:

- تأثیر1: توضیح مختصر
- تأثیر2: توضیح مختصر

محدودیت های شواهد چه بودند؟

مخاطبان طرح پژوهشی: رسانه و مردم
آیا این خبر می تواند از نظر سیاسی، فرهنگی، بهداشتی، ارزش های دینی و قوانین سازمان غذا و دارو، تبعاتی داشته باشد؟

بله

در صورتی که این طرح منتج به مقاله شده است لینک مقاله درج شود:
https://www.opastpublishers.com/peer-review/predictive-and-preventive-factors-related-to-of-head-lice-infestations-among-teenage-girls-student-application-of-health-7318.html

ایمیل ارتباطی و تلفن مجری اصلی طرح:
هادی علیزاده سیوکی 09155319756

alizadeh1@thums.ac.ir

منابع و مراجع :4 مرجع اصلی استفاده شده در طرح تحقیقاتی مورد نظر را ذکر نمایید

1. Al Bashtawy M, Hasna F. Pediculosis capitis among primary-school children in Mafraq Governorate, Jordan. 2012.
2. Falagas ME, Matthaiou DK, Rafailidis PI, Panos G, Pappas G. Worldwide prevalence of head lice. 2008.
3. Hosseini S, Rajabzadeh R, Shoraka V, Avaznia A, Shoraka H. Prevalence of pediculosis and its related factors among primary school students in Maneh-Semelghan district. Journal of North Khorasan University of Medical Sciences. 2014;6(1):43-9.
4. Motevalli Haghi S, Rafieiad J, Hosseni M. Epidemiology of Pediculosis and Its Associated Risk Factors in Primary-School Children of Sari, Mazandaran Province, in 2012-2013. Journal of Health. 2014;4(4):339-48.
5. Gholamiparizad E AM. Infestation with head lice and its influencing factors in primary school Shhravlam. J Avlam Uni Med Sci. 2002:30(29).
6. Zareban E BM, Moadi M ,Mehriofard H, Ghafari H. Evaluation of an educational program to reduce the prevalence of head lice among female students in elementary schools in Zabl. J Byrjnd Uni Med Sci. 2007:13(1):13.
7. Kassiri H, Farhadinejad R, Lotfi M. PHARMACEUTICAL SCIENCES.
8. Gholamnia Shirvani Z, Shokravi A, Ardestani M. Effect of designed health education program on knowledge, attitude, practice and the rate Pediculosis Capitis in female primary school students in Chabahar city. journal of shahrekord university of medical sciences. 2011;13(3).
9. Farzinnia B, Hanafibejad A, Raeiskarami S, Jafari T. Epidemiology of head lice in primary school girls in Qom, 2002. Hormozgan University of Medical Sciences. 2004;2(8):103-8.
10. Speare R, Cahill C, Thomas G. Head lice on pillows, and strategies to make a small risk even less. International journal of dermatology. 2003;42(8):626-9.
11. Kasiri H. Epidemiology of pediculus humanus capitis infestation and effective factors in elementary schools of girls Ahvaz city, 2005. Iran J Infect Dis Trop Med. 2009;14(2):41-5.
12. Mumcuoglu KY. Prevention and treatment of head lice in children. Pediatric Drugs. 1999;1(3):211-8.
13. Guvenc G, Akyuz A, Acikel CH. Health belief model scale for cervical cancer and Pap smear test: psychometric testing. Journal of advanced nursing. 2011;67(2):428-37.
14. Mazloomi Mahmoodabad S, Roohani Tanekaboni N. Survey of some related factors to oral health in high school female students in Yazd, on the basis of health behavior model (HBM). Journal of Birjand university of medical sciences. 2008;15(3):40-7.
15. NAJIMI A, ALIDOUSTI M, MOAZEMI GA. A survey on preventive behaviors of high school students about Influenza A based on health belief model in Shahrekord, Iran. 2010.
16. Sajadi Hazaveh M, Shamsi M. Assesment of mothers' behavior about prevention of febrile seizure in children in Arak city: application of the health belief model. Journal of Jahrom University of Medical Sciences. 2011;9(2):34-40.
17. Namdar A, Bigizadeh S, Naghizadeh MM. Measuring Health Belief Model components in adopting preventive behaviors of cervical cancer. Journal of Fasa University of Medical Sciences. 2012;2(1):34-44.
18. MorowatiSharifabad M, EbrahimZadeh M, Fazeli F, Dehghani A, Neshati T. Study of Pediculus capitis prevalence in primary school children and its preventive behaviors determinants based on Health Belief Model in Their Mothers in Hashtgerd, 2012. Tolooebehdasht. 2016;14(6):198-207.
19. Moshki M, Mojadam M, Alaviieh FZ. Preventive Behaviors of Female Elementary Students in regard to Pediculosis Infestation based on Health Belief Model (HBM). Journal of health and development. 2014;3(3):269-81.
20. Daneshvar S, Aivazi AA, Naghizadeh MM, Ghazanfari Z. Factors Associated with Preventive Behaviors of Pediculosis Infection among Elementary School Girl Students in Eyvan: An Application of the Health Belief Model. J Educ Community Health.0-.
21. GOODARZVAND CP, ANOOSHEH M, KAZEMNEJAD A. THE EFFECTIVENESS OF EDUCATING MOTHERS ON PREVENTIVE BEHAVIORS OF PEDICULOSIS ON MORBIDITY RATE OF THEIR DAUGHTERS. 2017.
22. Yingklang M, Senathong C, Haonon O, Dangtakot R, Pinlaor P, Sota C, et al. Effect of a health education program on reduction of pediculosis in school girls at Amphoe Muang, Khon Kaen Province, Thailand. PloS one. 2018;13(6):e0198599.

23. Mainbolagh BL, Rakhshani F, Zareban I, Montazerifar F, Sivaki HA, Parvizi Z. The effect of peer education based on health belief model on nutrition behaviors in primary school boys. Journal of Research & Health. 2012;2(2):214-25.